



PHR

S
Y
N
O
P
S
E
S

MORRIS, LEO (Public Health Service): Further analysis of national participation in the poliomyelitis vaccination program, 1955-61. Public Health Reports, Vol. 79, June 1964, pp. 469-480.

Since 1957 the Public Health Service has sponsored annual surveys to determine nationwide participation in the inactivated poliomyelitis vaccination program which began in April 1955. These vaccination surveys were conducted in 5 successive years by the Bureau of the Census as supplements to the monthly Current Population Surveys.

The proportion of the population adequately immunized has progressively increased since the licensure of Salk vaccine in April 1955. By September 1961, 77 percent of the population under 40 years of age had received at least one inoculation, about 67 percent had received three or more inoculations, and 40 percent had received the fourth or booster inoculation. Approximately 60 percent of school-age children had four or more doses of inactivated poliomyelitis vaccine by September 1961, but less than half of the preschool children and young adults and

less than one-quarter of the men had reached this level.

The response of preschool children, adult males, and lower socioeconomic groups has been well below the participation of school-age children, adult females, and higher socioeconomic groups respectively. Since the advent of the inactivated poliomyelitis vaccine in 1955, poliomyelitis has attacked these poorly immunized groups with greater frequency.

Estimates of the effectiveness of three or more doses of inactivated poliomyelitis vaccine, as studied in four recent epidemics, have ranged from 77 to 82 percent. The sharp drop in poliomyelitis incidence since 1955 can largely be attributed to the use of the inactivated vaccine. That the decrease in paralytic poliomyelitis has not been greater may be attributed to the population groups who have remained poorly immunized.

BROWN, WILLIAM J. (Communicable Disease Center), DONOHUE, JAMES F., and PRICE, ELEANOR V.: Evaluation of RPR card test for syphilis screening in field investigations. Public Health Reports, Vol. 79, June 1964, pp. 496-500.

After a 2-day training course at the Venereal Disease Research Laboratory, 28 nontechnical venereal disease investigators performed the rapid plasma reagin card test on more than 3,900 persons. The VDRL slide test was also performed on approximately 2,800 of these persons by local laboratory personnel. Agreement between the two tests was 93 per-

cent; where discrepancies occurred, positivity was usually in the low range.

It was the consensus of the investigators that the RPR card test is impractical for confidential investigations conducted outside the clinic. The tests is a valuable tool, however, for improving clinic efficiency, since test results are immediately available.

HENDERSON, MAUREEN (University of Maryland School of Medicine), GOLDSTEIN, HYMAN, ROGOT, EUGENE, GOLDBERG, IRVING D., and ENTWISLE, GEORGE: Perinatal factors associated with epilepsy in Negro children. Public Health Reports, Vol. 79, June 1964, pp. 501-509.

A total of 299 Negro epileptic children born since 1950 in Maryland or Washington, D.C., were studied retrospectively. Each index case was matched, from birth certificates, with a control case for race, sex, plurality, birth weight, hospital of delivery, age of mother, and date of delivery. Children with diagnoses of mental retardation or cerebral palsy were eliminated from the case group.


Review of the hospital prenatal and delivery records showed that the pregnancies of the 283 mothers of children in the case group were similar to that of mothers of the control group in the fre-

quency of hemorrhagic, toxemic, and other medical complications. Although not statistically significant, the mothers of epileptic children had more mechanical complications of delivery. A somewhat greater proportion of mothers of the case children than mothers of controls had at least one previous pregnancy loss (abortion or stillbirth), and a greater proportion of the epileptic children were delivered by cesarean section.

In addition, comparison with a second matched group showed that children with epilepsy were not characterized by low birth weight.

CONTENTS *continued*

| | <i>Page</i> |
|---|-------------|
| Differences in glucose determinations obtained from plasma or whole blood. | 515 |
| <i>Glen W. McDonald, Gail F. Fisher, and Clinton E. Burnham</i> | |
| An evaluation of the role of <i>Phormia regina</i> in transmission of eastern encephalitis. | 522 |
| <i>A. T. C. Bourke</i> | |
| Factors influencing CVA patients to seek rehabilitative care. | 525 |
| <i>Charles M. Wylie</i> | |
| The nurse's role in rehabilitation of the child with rheumatic fever. | 533 |
| <i>Lillian S. Dick and Marie D. Grant</i> | |
| Coordinating a community study of environmental health. | 537 |
| <i>Walter C. Kane</i> | |
| Sanitary landfill operations in New York State. | 543 |
| <i>Salvatore Pagano</i> | |
| The grants associates program. | 549 |
| Short reports and announcements: | |
| Study on drug-treated schizophrenics. | 467 |
| Resources for mentally ill children. | 480 |
| Study on rehabilitation of drug addicts. | 487 |
| New chief for Division of Hospitals. | 500 |
| Sharing project and activity information. | 521 |
| Studies on radioactivity in children's diets. | 532 |
| Program notes. | 542 |
| Federal publications. | 551 |



MANAGING DIRECTOR

J. STEWART HUNTER, M.A.
*Assistant to the Surgeon General
 for Information
 Office of Information and Publications*

BOARD OF EDITORS

GEORGE ST. J. PERROTT
Chairman

BERNARD V. DRYER, M.D.
 DONALD J. GALAGAN, D.D.S., M.P.H.
 LEO J. GEHRIG, M.D.
 JAMES HUNDLEY, M.D.
 ROSCOE P. KANDLE, M.D., M.P.H.
 F. ELLIS KELSEY, PH.D.
 LUCILE P. LEONE, R.N., M.A.
 DAVID LITTAUER, M.D.
 MARGARET F. SHACKELFORD, M.S.
 ELLIS D. SOX, M.D.
 JAMES WATT, M.D., DR.P.H.

MARCUS ROSENBLUM
Secretary to the Board

STAFF

Keith Kost, M.P.H. *Editor*
 Winona Carson *Managing Editor*

Address correspondence to Editor, Public Health Reports, Public Health Service, Department of Health, Education, and Welfare, Washington, D.C., 20201.

Opinions expressed are the authors' and do not necessarily reflect the views of Public Health Reports or the Public Health Service. Trade names are used for identification only and do not represent an endorsement by the Public Health Service.

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE

ANTHONY J. CELEBREZZE, *Secretary*

PUBLIC HEALTH SERVICE

LUTHER L. TERRY, *Surgeon General*

PHR

S
Y
N
O
P
S
E
S

MYERBURG, ROBERT J. (Public Health Service), **JABLON, JAMES M.**, **MAZ-ZARELLA, JOHN A.**, and **SASLAW, MILTON S.**: *Evaluation of FA and conventional techniques for identifying group A beta hemolytic streptococci.* *Public Health Reports, Vol. 79, June 1964, pp. 510-514.*

Ten bacteriological and immunofluorescent procedures for the identification of group A beta hemolytic streptococci were evaluated to determine the most rapid and accurate technique for mass epidemiological studies. Five fluorescent antibody technique modifications, a pour plate technique, three streak plate modifications, and the conventional microprecipitin grouping technique were tested.

Group A beta hemolytic streptococci were recovered from 177 of a total of 1,115 throat cultures obtained from children with respiratory illnesses. Best results were obtained when a colony was picked from a pour plate after 18 hours' incubation at 37° C. and then identified by

fluorescent antibody microscopy. No group A strains were missed; 3 of 956 cultures were falsely identified. Other fluorescent antibody methods met with varying degrees of success, but none approached the accuracy of the above method.

The three streak plate techniques were unsuccessful. The pour plate technique was excellent for indication of beta hemolysis; only one group A strain was missed, but many non-group A organisms also produced beta hemolysis and were thus indistinguishable.

The conventional microprecipitin grouping technique was used as the control method for this study.

McDONALD, GLEN W. (Public Health Service), **FISHER, GAIL F.**, and **BURNHAM, CLINTON E.**: *Differences in glucose determinations obtained from plasma or whole blood.* *Public Health Reports, Vol. 79, June 1964, pp. 515-521.*

Glucose determinations obtained on the AutoAnalyzer from plasma are consistently higher than those obtained from whole blood. The difference is not absolute but tends to increase as the level of the glucose increases. Since there is a close relationship between plasma and whole blood glucose values obtained from the same samples, it is possible to set critical levels which correspond. Such levels are presented for values used frequently in diabetes screening programs

or in the interpretation of the standard 100 gm. oral glucose tolerance test. The relationship of glucose values obtained from whole blood and plasma is sufficiently precise that interpretation of data based on whole blood or based on plasma is equally valid as long as adjustments for differences in levels are made. Data are presented which permit the estimation of equivalent values for levels other than those presented.

The nature of a paper, not its importance or significance, determines whether a synopsis is printed. See "Information for Contributors" on last page.

Information for Contributors

PUBLIC HEALTH REPORTS welcomes from any source all contributions of value to public health.

Most of the readers of *Public Health Reports* are practicing public health officials. About 10 percent of the monthly circulation of *Public Health Reports* goes overseas. About half of the domestic circulation goes to Federal, State, and local government agencies concerned with health and related interests. A quarter goes to institutions accredited for teaching in health and related fields, to teachers, and to libraries. The journal also reaches research institutions, hospitals, and professional and voluntary public health organizations.

Tearsheets. In lieu of reprints, senior authors are provided with 50 to 100 sets of tearsheets after publication. Associate authors receive a smaller number.

Manuscript review. Manuscripts submitted for publication are reviewed by technical experts, and authors are given the benefit of their comments before type is set. Authors also receive edited typescripts for approval and are given the opportunity to correct galley proofs. Authors are responsible for the accuracy and validity of all material, including tables, charts, and references. Special editorial assistance in preparing or revising manuscripts is available on request, to the limit of staff resources.

Manuscripts are reviewed with the understanding that they have not been committed for publication elsewhere. Appropriate information should be provided if a paper has been given or is prepared for presentation at a meeting.

Manuscript form. Authors will facilitate review and publication if they submit an original and three carbon copies of their manuscripts. All copy should be typed double spaced, and each page should end with a completed paragraph. Of course, several paragraphs may appear on a typed page.

References should be given in the style used by *Public Health Reports*.

Footnotes should be worked into the text or offered as supplemental items.

Authors are expected to recognize scientific contributions by those who have assisted in their papers only if such contributions warrant mention in the text or in the paragraph identifying the authors. It is not the policy of *Public Health Reports* to publish "acknowledgments."

Synopses. To facilitate secondary publication, *Public Health Reports* publishes synopses of selected papers, principally research studies. Authors are requested to submit with appropriate papers a synopsis of not more than 200 words. The staff will supply on request information on preparation of synopses.

Secondary publication. Secondary publication of articles in *Public Health Reports* is provided in various abstracting journals. Articles are also indexed in the annual Cumulated Index Medicus (American Medical Association), the monthly Index Medicus (National Library of Medicine), the Engineering Index, the Hospital Literature Index, and the biannual supplements to the Cumulative Index to Nursing Literature.

Bound copies. Librarians and others should preserve their copies for binding, as the Public Health Service does not supply bound copies. Indexes are published each year in the December issue.

PUBLIC HEALTH MONOGRAPHS, edited and issued by *Public Health Reports*, must be submitted through constituent agencies of the Department of Health, Education, and Welfare.

Most Public Health Monographs are placed on sale by the Superintendent of Documents; series subscriptions are not available. Monographs are not included in subscriptions to *Public Health Reports*.

Address correspondence on editorial matters to:
Editor, *Public Health Reports*, Public Health Service, U.S. Department of Health, Education, and Welfare, Washington, D.C., 20201.